









LISTING OF CLAIMS

1. (Currently Amended) An ionic liquid of the general formula



wherein:

K^+ is a cation selected from:




wherein

R^1 to R^6 are identical or different and are each individually

- H,
- a halogen,

- an alkyl radical (C_1 to C_8), which is unsubstituted, or which is partially or fully substituted by F, Cl, $N(C_nF_{(2n+1-x)}H_x)_2$, $O(C_nF_{(2n+1-x)}H_x)$, $SO_2(C_nF_{(2n+1-x)}H_x)$ or $C_nF_{(2n+1-x)}H_x$ wherein $1 < n < 6$ and $0 < x \leq 13$
- a phenyl radical which is unsubstituted or which is partially or fully substituted by F, Cl, $N(C_nF_{(2n+1-x)}H_x)_2$, $O(C_nF_{(2n+1-x)}H_x)$, $SO_2(C_nF_{(2n+1-x)}H_x)$ or $C_nF_{(2n+1-x)}H_x$ wherein $1 < n < 6$ and $0 < x \leq 13$, or
- one or more pairs of adjacent R^1 to R^6 can also be an alkylene or alkenylene radical and having up to 8 C atoms, wherein the radical is unsubstituted or partially or fully substituted by halogen, $N(C_nF_{(2n+1-x)}H_x)_2$, $O(C_nF_{(2n+1-x)}H_x)$, $SO_2(C_nF_{(2n+1-x)}H_x)$ or $C_nF_{(2n+1-x)}H_x$ wherein $1 < n < 6$ and $0 < x \leq 13$

wherein A^+ is an anion selected from



wherein

$0 \leq n, m, o, p \leq 4$, and $m+n+o+p=4$, and

R^7 to R^{10} are different or identical and are each, individually:

an aromatic ring selected from a phenyl, ~~naphthyl~~, anthracenyl and phenanthrenyl ring, which is unsubstituted, or which is monosubstituted or polysubstituted by $C_nF_{(2n+1-x)}H_x$, wherein $1 < n < 6$ and $0 < x \leq 13$, or halogen,

an aromatic heterocyclic ring selected from a pyridyl, pyrazyl and pyrimidyl ring, which is unsubstituted, or which is mono-substituted or polysubstituted by $C_nF_{(2n+1-x)}H_x$, wherein $1 < n < 6$ and $0 < x \leq 13$, or halogen,

or

an alkyl radical (C_1 to C_8), which is unsubstituted, or which is partially or fully substituted by F, Cl, $N(C_nF_{(2n+1-x)}H_x)_2$, $O(C_nF_{(2n+1-x)}H_x)$, $SO_2(C_nF_{(2n+1-x)}H_x)$, or $C_nF_{(2n+1-x)}H_x$, wherein $1 < n < 6$ and $0 < x \leq 13$,

and wherein one or more pairs of R^7 to R^{10} can also form

an aromatic ring selected from a ~~phenylene, naphthylene,~~
anthracenylene and phenanthrenylene ring, which is unsubstituted
or an aromatic ring selected from a phenylene, naphthylene,
anthracenylene and phenanthrenylene ring which is
monosubstituted or polysubstituted by $C_nF_{(2n+1-x)}H_x$, wherein $1 < n < 6$
and $0 < x \leq 13$, or halogen,

an aromatic heterocyclic ring selected from a pyridylene,
pyrazylene and pyrimidylene ring, which is unsubstituted, or which
is mono-substituted or polysubstituted by $C_nF_{(2n+1-x)}H_x$, wherein
 $1 < n < 6$ and $0 < x \leq 13$, or halogen,

or

an alkylene or alkenylene radical having up to 8 C atoms and which
is unsubstituted or which is partially or fully substituted by halogen,
 $N(C_nF_{(2n+1-x)}H_x)_2$, $O(C_nF_{(2n+1-x)}H_x)$, $SO_2(C_nF_{(2n+1-x)}H_x)$ or $C_nF_{(2n+1-x)}H_x$
wherein $1 < n < 6$ and $0 < x \leq 13$

or OR^7 to OR^{10} , individually or together,

are an aromatic having 6 to 14 C atoms and which is a ~~carboxyl,~~
dicarboxyl, oxysulfonyl or oxycarbonyl radical, which is
unsubstituted, or which is partially or fully substituted by F, Cl,
 $N(C_nF_{(2n+1-x)}H_x)_2$, $O(C_nF_{(2n+1-x)}H_x)$, $SO_2(C_nF_{(2n+1-x)}H_x)$ or $C_nF_{(2n+1-x)}H_x$,
wherein $1 < n < 6$ and $0 < x \leq 13$

or

are aliphatic having 1 to 6 C atoms and which is a carboxyl, dicarboxyl, oxysulfonyl or oxycarbonyl radical, which is unsubstituted, or which is partially or fully substituted by F, Cl, $N(C_nF_{(2n+1-x)}H_x)_2$, $O(C_nF_{(2n+1-x)}H_x)$, $SO_2(C_nF_{(2n+1-x)}H_x)$ or $C_nF_{(2n+1-x)}H_x$ wherein $1 < n < 6$ and $0 < x \leq 13$.

2. **(original claim)** An ionic liquid according to claim 1, wherein at least one of R^1 to R^6 of the cation is an alkyl radical which is unsubstituted or partially or fully substituted by F, Cl, $N(C_nF_{(2n+1-x)}H_x)_2$, $O(C_nF_{(2n+1-x)}H_x)$, $SO_2(C_nF_{(2n+1-x)}H_x)$ or $C_nF_{(2n+1-x)}H_x$ wherein $1 < n < 6$ and $0 < x \leq 13$.

3 **(original claim)** An ionic liquid according to claim 1, wherein at least one of R^1 to R^6 of the cation is a phenyl radical which is unsubstituted or partially or fully substituted by F, Cl, $N(C_nF_{(2n+1-x)}H_x)_2$, $O(C_nF_{(2n+1-x)}H_x)$, $SO_2(C_nF_{(2n+1-x)}H_x)$ or $C_nF_{(2n+1-x)}H_x$ wherein $1 < n < 6$ and $0 < x \leq 13$.

4. **(original claim)** An ionic liquid according to claim 1, wherein at least a pair of R^1 to R^6 of the cation is an alkylene or alkenylene radical which is unsubstituted or partially or fully substituted by halogen, $N(C_nF_{(2n+1-x)}H_x)_2$, $O(C_nF_{(2n+1-x)}H_x)$, $SO_2(C_nF_{(2n+1-x)}H_x)$ or $C_nF_{(2n+1-x)}H_x$ wherein $1 < n < 6$ and $0 < x \leq 13$.

5. **(original claim)** An ionic liquid according to claim 1, wherein at least one of R^7 to R^{10} of the anion is an alkyl radical which is unsubstituted or partially or fully substituted by F, Cl, $N(C_nF_{(2n+1-x)}H_x)_2$, $O(C_nF_{(2n+1-x)}H_x)$, $SO_2(C_nF_{(2n+1-x)}H_x)$, or $C_nF_{(2n+1-x)}H_x$ wherein $1 < n < 6$ and $0 < x \leq 13$.

6. **(original claim)** An ionic liquid according to claim 1, wherein at least one pair of R^7 to R^{10} of the anion is an alkylene or alkenylene radical which is unsubstituted or partially or fully substituted by a halogen, $N(C_nF_{(2n+1-x)}H_x)_2$, $O(C_nF_{(2n+1-x)}H_x)$, $SO_2(C_nF_{(2n+1-x)}H_x)$ or $C_nF_{(2n+1-x)}H_x$ wherein $1 < n < 6$ and $0 < x \leq 13$.

7. **(currently amended)** An ionic liquid according to claim 1, wherein at least one of R^7 to R^{10} of the anion is an aromatic ring selected from a phenyl, ~~naphthyl~~, anthracenyl and phenanthrenyl ring, which is unsubstituted, or which is monosubstituted or polysubstituted by $C_nF_{(2n+1-x)}H_x$ wherein $1 < n < 6$ and $0 < x \leq 13$, or by a halogen.

8. **(currently amended)** An ionic liquid according to claim 1, wherein at least one of R^7 to R^{10} of the anion is an aromatic heterocyclic ring selected from a pyridyl, pyrazyl and pyrimidyl ring, which is unsubstituted, or which is monosubstituted or polysubstituted by $C_nF_{(2n+1-x)}H_x$ wherein $1 < n < 6$ and $0 < x \leq 13$, or ~~by a halogen~~ (F, Cl or Br).

9. **(currently amended)** An ionic liquid according to claim 1, wherein at least one pair of R^7 to R^{10} of the anion is an aromatic ring selected from a ~~phenylene, naphthylene~~, anthracenylene and phenanthrenylene ring, which is unsubstituted or a phenylene, naphthylene, anthracenylene and phenanthrenylene ring, which is monosubstituted or polysubstituted by $C_nF_{(2n+1-x)}H_x$ wherein $1 < n < 6$ and $0 < x \leq 13$, or halogen.

10. **(original claim)** An ionic liquid according to claim 1, wherein at least one pair of R^7 to R^{10} of the anion is an aromatic heterocyclic ring selected from a pyridylene, pyrazylene and pyrimidylene ring, which is unsubstituted, or which is mono-substituted or polysubstituted by $C_nF_{(2n+1-x)}H_x$ wherein $1 < n < 6$ and $0 < x \leq 13$, or by halogen.

11. **(withdrawn)** An electrochemical cell comprising a cathode, an anode, a separator, and the ionic liquid of claim 1.

12. **(withdrawn)** A supercapacitor comprised of at least a pair of electrodes, a separator, and the ionic liquid of claim 1.

13. **(withdrawn)** An electrolyte composition comprising an ionic liquid of claim 1 and an aprotic solvent.

14. **(withdrawn)** An electrolyte composition comprising an ionic liquid of claim 1 and a conductive salt.

15. **(original claim)** A method for making an ionic liquid according to claim 1, comprising reacting a chloride salt of the formula K^+Cl^- with a lithium salt of the formula Li^+A^- within an aprotic solvent.

16. **(currently amended)** A compound ~~according to claim 1, wherein~~
~~said compound is~~ selected from :

1-ethyl-3-methylimidazolium bis [1,2-benzenediolato-O,O'] borate,

1-ethyl-3-methylimidazolium bis[oxalato]borate, ~~or and~~

1-ethyl-3-methylimidazolium bis[salicylato]borate.

17. (currently amended) A compound according to claim 4 16, wherein said compound is:

1-ethyl-3-methylimidazolium bis [1,2-benzenediolato-O,O'] borate.

18. (new) A compound according to claim 1, wherein A⁺ is

bis[oxalato]borate,

or

bis[salicylato]borate.